

Randall-Sundrum with AdS(7)

Ruoyu Bao

*Enrico Fermi Institute and Department of Physics,
University of Chicago, Chicago, IL 60637, USA*

Joseph Lykken

*Theoretical Physics Department,
Fermi National Accelerator Laboratory,
P.O. Box 500, Batavia, IL 60510 USA*

In the same sense that AdS_5 warped geometries arise naturally from Type IIB string theory with stacks of $D3$ branes, AdS_7 warped geometries arise naturally from M theory with stacks of $M5$ branes. We compactify two spatial dimensions of AdS_7 to get $AdS_5 \times \Sigma^2$, where the metric for Σ^2 inherits the same warp factor as appears in the AdS_5 . We analyze the 5d spectrum in detail for the case of a bulk scalar or a graviton in $AdS_5 \times T^2$, in a setup which mimics the first Randall-Sundrum model. The results display novel features which might be observed in experiments at the CERN Large Hadron Collider. For example, we obtain TeV scale string winding states without lowering the string scale. This is due to the *double warping* which is a generic feature of winding states along compactified AdS directions. Experimental verification of these signatures of AdS_7 could be interpreted as direct evidence for M theory.

PACS numbers: 11.25.Mj, 11.25.Uv, 11.25.Yb, 11.10.Kk

FERMILAB-PUB-05-381-T

INTRODUCTION

The five-dimensional warped models of Randall and Sundrum [1] can be thought of as stripped down effective field theory approximations to the near horizon geometry [2] of a stack of N parallel $D3$ branes in Type IIB string theory. This near horizon geometry is $AdS_5 \times S^5$, where the Anti-de Sitter radius R_{AdS} and the radius of the sphere are equal, scaling like $N^{1/4}$. Thus for large N the AdS physics involves an energy scale that is parametrically lower than the string (or Planck) scale. So does the physics on S^5 , but we also expect this physics to be drastically altered as one breaks supersymmetry, and otherwise attempt to make the compactification more realistic. Since in any event the sphere S^5 is not warped, it seems reasonable to factor out this physics and examine low energy effective 5d field theory on AdS_5 . In the Randall-Sundrum (RS) approach, a phenomenological 3-brane called the Planck brane is added to truncate AdS_5 in a way that excludes the AdS boundary. From the point of view of string theory, this is a boundary condition that replaces the original matching of the near horizon geometry to $M_5 \times S^5$; in the AdS/CFT dual language, the Planck brane is a particular choice of UV cutoff for the 4d CFT coupled to 4d gravity. The first Randall-Sundrum model (RS1) also introduces a second brane, called the TeV brane. This brane truncates the AdS geometry in the other direction, excluding now the AdS horizon. This is a wise idea for a low energy effective field theory approach, since close to the horizon the field theory cutoff necessarily shrinks to zero and stringy effects become important.

In this paper we examine an analogous story which

starts with eleven dimensional M theory. The low energy supergravity theory has a solution [2] corresponding to a stack of N parallel $M5$ branes. These $M5$ branes are just the magnetic sources for the antisymmetric tensor gauge field A_{MNP} of 11d supergravity, which also has electric sources called $M2$ branes. We extract the near horizon geometry of this solution for (moderately) large N . This geometry is $AdS_7 \times S^4$, where $R_{AdS} = 2(\pi N)^{1/3} \ell_p$, with ℓ_p the Planck length, in the convention where the $M2$ branes have inverse tension $(2\pi)^2 \ell_p^3$.

As motivated above we will discard the unwarped sphere S^4 and consider low energy effective 7d field theories built on AdS_7 :

$$ds^2 = a^2(y) (\eta_{\mu\nu} dx^\mu dx^\nu + \delta_{ij} dx^i dx^j) + dy^2, \quad (1)$$

where we have introduced the warp factor $a(y) = \exp(-ky)$, $k = 1/R_{AdS}$. Our metric signature convention is $(- + + + +)$. As in RS, we have to tune the brane tensions to obtain flat 6d slices. Note that AdS_7 setups are very restricted by the requirement of anomaly cancellation in the 6d boundary gauge theories [3].

AdS₅ WITH EXTRA WARPINGS

To make contact with a Randall-Sundrum type construction, we now compactify two spatial dimensions of the AdS_7 . Thus

$$AdS_7 \rightarrow AdS_5 \times \Sigma^2.$$

The simplest choice for Σ^2 is a torus T^2 , since in this case we do not need any additional sources in the Einstein

equations to get a consistent background solution. The metric becomes:

$$ds^2 = a^2(y)\eta_{\mu\nu}dx^\mu dx^\nu + a^2(y)R^2(d\theta_1^2 + d\theta_2^2) + dy^2, \quad (2)$$

where for simplicity we assume a common radius parameter R . Note that the measured radius of the torus includes a warp factor.

Almost as simple is a compactification on a sphere S^2 . Since the sphere has curvature, this requires an additional bulk source. As discussed for example in [4], two bulk scalar fields with a simple potential allows a solution of the equations of motion with a “hedgehog” profile for the scalars.

Any 7d bulk field can be expanded in a complete orthonormal basis of Kaluza-Klein (KK) modes for Σ^2 . These are just sines and cosines for T^2 , and spherical harmonics for S^2 . Substituting the KK expansion into the 7d action, we can then integrate over Σ^2 to obtain an effective 5d action.

We will do this explicitly for the case of a 7d bulk scalar on $AdS_5 \times T^2$, and then extend the results to discuss the 7d graviton. For the torus with radius R the squared KK momenta are

$$g^{ij}p_i p_j = \frac{n_1^2 + n_2^2}{a^2 R^2} \equiv \frac{\vec{n}^2}{a^2 R^2}, \quad (3)$$

where n_1, n_2 are integers.

In the underlying theory there will in general be additional bulk modes arising from $M2$ or $M5$ branes wrapping one or more cycles of the torus. These will be particle-like modes if the other dimensions of the branes are already wrapped around compact cycles of S^4 , or whatever replaces S^4 in a realistic model. For example, an $M5$ brane wrapped on S^4 is a baryonic string [5], which can in turn wind around cycles of the torus. Such modes can be labelled by their winding numbers, which form a two-vector of integers $m^i, i = 1, 2$.

Whereas the KK mode contributions to the world-sheet Hamiltonian of the string are proportional to $\ell_p^2 g^{ij} p_i p_j$, the winding modes contribute

$$\frac{R^2}{\ell_p^2} g_{ij} m^i m^j = \frac{R^2}{\ell_p^2} a^2 \vec{m}^2. \quad (4)$$

Note that the parameter ℓ_p above will in general differ from the ℓ_p introduced above (1), by a factor of order one which depends upon how strings emerge from the original 11d M theory.

We can write down the equations of motion for the bulk scalar in the 5d effective theory, expanded into KK and winding modes $\phi_{\vec{n}\vec{m}}(x^\mu, y)$:

$$\left[\frac{1}{a^4} \frac{d}{dy} a^6 \frac{d}{dy} - p^2 + \frac{\vec{n}^2}{R^2} - a^2 m_b^2 - a^4 \frac{\vec{m}^2 R^2}{\ell_p^4} \right] \phi_{\vec{n}\vec{m}} = 0, \quad (5)$$

where m_b is an explicit bulk mass from the 7d theory.

The usual warping of Randall-Sundrum is visible in (5) as the fact that the bulk mass-squared term is multiplied by $a^2(y)$ relative to the 4d momentum-squared $p^2 = p_\mu p^\mu$. We observe as well two new kinds of behavior:

- The KK mode mass-squared contributions from the torus have no warping relative to the 4d momentum-squared $p_\mu p^\mu$.
- The winding mode mass-squared contributions have double warping compared to the ordinary bulk mass term, *i.e.*, they are multiplied by $a^4(y)$ relative to the 4d momentum-squared $p_\mu p^\mu$.

Now we change variables to the conformal coordinate $z = 1/ka(y)$, suppress indices, and perform the useful rescaling $\phi(z) = k^3 z^3 h(z)$. The equation of motion becomes:

$$z^2 h'' + z h' + \left[(\mu^2 - \frac{\vec{n}^2}{R^2}) z^2 - \nu^2 - \frac{\vec{m}^2 R^2}{k^4 \ell_p^4} \frac{1}{z^2} \right] h = 0, \quad (6)$$

where $\nu^2 = 9 + m_b^2/k^2$, and we are putting the 4d momenta on-shell, with $\mu^2 = -p_\mu p^\mu$ denoting the physical 4d mass. It is the mass spectrum of μ that we want to determine from an analysis of the solutions of (6). For $\vec{n}^2 = \vec{m}^2 = 0$, this is the equation of motion for a 5d massive bulk scalar in AdS_5 , as first analyzed in [6]. In general (6) is equivalent to the Mathieu equation [7].

5D KALUZA-KLEIN MODE ANALYSIS

We can take (6) as the starting point for a 5d Kaluza-Klein mode analysis in AdS_5 . To get the analog of scalars in a Randall-Sundrum model, we can impose Neumann boundary conditions at the location of one or two flat branes. These branes have co-dimension one, *i.e.* they are 5-branes, located at $z = 1/k$ and $z = 1/T$. The parameter T is assumed to be of order the TeV scale.

To analyze graviton modes, we use the fact that in *e.g.* harmonic gauge each component of the graviton obeys (6). The appropriate boundary conditions for the graviton modes, after tuning the brane tensions to their conventional RS values, are:

$$[zh'(z) + 3h(z)]_{z=1/k} = [zh'(z) + 3h(z)]_{z=1/T} = 0. \quad (7)$$

This means that the graviton mode solutions are identical to the $m_b = 0$ scalar solutions, up to an overall factor.

Let us start with the simple case where $\vec{m} = 0$. The solution to (6) depends upon the sign of $\mu^2 - \vec{n}^2/R^2$. There are no solutions for $\mu^2 - \vec{n}^2/R^2 < 0$. For $\mu^2 - \vec{n}^2/R^2 > 0$ the solution is

$$\phi(z) = \frac{k^3 z^3}{N} \left[J_\nu \left(z \sqrt{\mu^2 - \frac{\vec{n}^2}{R^2}} \right) + b Y_\nu \left(z \sqrt{\mu^2 - \frac{\vec{n}^2}{R^2}} \right) \right]. \quad (8)$$

If we only consider the case of light modes where $\mu^2 - \frac{\tilde{n}^2}{R^2} \ll k^2$, then the boundary conditions determine b and yield

$$3J_\nu(x) + xJ'_\nu(x) = 0, \quad (9)$$

where $xT = \sqrt{\mu^2 - \frac{\tilde{n}^2}{R^2}}$. The roots of (9) determine the mass spectrum. These solutions represent towers of toroidal KK modes on top of the TeV-spaced massive warped KK modes of Randall-Sundrum. The RS spectrum itself is also modified, since *e.g.*, for $m_b = 0$, $\tilde{n} = 0$ the relevant Bessel functions are $J_3(\mu z)$ and $Y_3(\mu z)$, rather than $J_2(\mu z)$ and $Y_2(\mu z)$ as in standard RS.

Since the toroidal KK modes are not warped, $R \gtrsim 1/T$ corresponds to compactifications large enough that we will see the toroidal KK excitations with spacings comparable to the Randall-Sundrum KK modes. In order not to reintroduce the hierarchy problem, we would then need some mechanism that naturally stabilizes R at inverse TeV values. By contrast, if our torus were compactifying flat rather than *AdS* directions, then $R \sim 1/k$ would give TeV-spaced KK excitations [8].

In the special case $\mu^2 - \tilde{n}^2/R^2 = 0$, the solution to the equation of motion is

$$\phi(z) = \frac{k^3 z^3}{N} (z^{-\nu} + bz^\nu). \quad (10)$$

Similarly, the boundary conditions give us

$$(3 - \nu)k^\nu + b(3 + \nu)k^{-\nu} = 0, \quad (11)$$

$$(3 - \nu)T^\nu + b(3 + \nu)T^{-\nu} = 0. \quad (12)$$

This set of equations has no solution except when $\nu = 3$, *i.e.*, when there is a scalar zero mode. The solutions then represent the tower of toroidal KK modes on top of the zero mode.

5D WINDING MODE ANALYSIS

The most interesting new feature of our analysis is the spectrum of winding modes. For simplicity, we will only consider the simplest case, with $\tilde{n} = 0$ and $m_b = 0$. Then (6) reduces to:

$$z^2 h'' + zh' + \left(\mu^2 z^2 - 9 - \frac{\tilde{R}^2}{z^2} \right) h = 0, \quad (13)$$

where we have introduced the notation $\tilde{R} = R|\tilde{m}|/k^2 \ell_p^2$. As usual in Randall-Sundrum setups, we will assume that $k\ell_p$ is less than 1 but not dramatically so, *e.g.*, $k\ell_p \simeq 0.1$.

First of all, we want to see if there exists a zero mode. Setting $\mu = 0$ the solution of (13) is

$$\phi(z) = \frac{k^3 z^3}{N} \left[I_\nu\left(\frac{\tilde{R}}{z}\right) + bK_\nu\left(\frac{\tilde{R}}{z}\right) \right]. \quad (14)$$

The boundary conditions give:

$$b = -\frac{3I_\nu(k\tilde{R}) - k\tilde{R}I'_\nu(k\tilde{R})}{3K_\nu(k\tilde{R}) - k\tilde{R}K'_\nu(k\tilde{R})} = [k \leftrightarrow T], \quad (15)$$

which cannot both be true at the same time unless $T = k$. Therefore there is no zero mode for the case with winding modes only.

For the massive winding modes, the solutions can be found numerically. Here we summarize the results.

$$R \sim 1/k$$

This corresponds to the smallest compactification radius, where we expect the winding modes to be the lightest. Like the KK modes, these should appear as additional excitations on top of the TeV-spaced warped KK modes of Randall-Sundrum. We find that the RS zero mode does not have any winding excitations. The winding excitations of the lowest lying massive RS modes have masses given by $\mu = \mu_1 - 0.40\tilde{R}^2 T^3$ and $\mu = \mu_2 + 1.46\tilde{R}^2 T^3$, where $\mu_1 = 5.13T$ and $\mu_2 = 8.42T$ are the masses of the first two massive RS modes. Notice that the mass splittings are very small, of order $10^4 \times T^3/k^2 \sim 10^{-12}$ eV.

$$R \sim 1/T$$

This corresponds to a large compactification radius, thus were it not for warping we would expect the winding excitations to be extremely heavy. However, because of the double warping of the winding term in (6), we find that in this case the winding excitations are actually of order T . Once again the RS zero mode does not have any winding excitations. The winding excitations of the lowest lying massive RS mode have masses as shown in Figure 1. Note that the spacing of the winding modes can be considerably less than T if R is small, *e.g.*, for $RT \sim 0.01$ and $k\ell_p \simeq 0.1$ the spacing is a fraction of T .

DOUBLE WARPING VERSUS T-DUALITY

Our results for the spectrum of KK and winding modes can be understood qualitatively as a combination of warping with the usual T-duality relation between the KK and winding spectra. For simplicity we will not distinguish between k and ℓ_p in this discussion.

When the compactification radius R is of order $1/T$, the toroidal KK modes are of order T , with no additional warping. The T-dual winding modes are of order k^2/T , multiplied by a double warp factor T^2/k^2 ; thus the winding modes are of order $k^2/T \times T^2/k^2 \sim T$, as reported above.




FIG. 1: Winding excitations of the lightest RS massive mode, for $\tilde{R}$ of order $1/T$.

When the compactification radius R is of order $1/k$, the toroidal KK modes are of order k , with no additional warping. The T-dual winding modes are of order k as well, but again multiplied by a double warp factor T^2/k^2 . The winding excitations are on top of the massive warped KK excitations, which are of order T . Since the masses add in quadrature, we expect the splittings of the winding mode excitations to be of order

$$\sqrt{T^2 + \frac{T^4}{k^2}} - T \sim \frac{T^3}{k^2}, \quad (16)$$

as reported above.

LHC DISCOVERY OPPORTUNITIES

The $AdS_7 \rightarrow AdS_5 \times \Sigma^2$ scenarios presented here are as well-motivated as the AdS_5 effective theories that are the basis for standard Randall-Sundrum. Both are stripped down versions of physics that could actually emerge from string theory in a robust way, albeit decorated with many stringy complications.

The spectrum of bulk graviton and scalar modes in our AdS_7 models can differ from standard RS1 in several ways. First of all, because we start with AdS_7 instead of AdS_5 , the spacing of the massive graviton modes is determined by the zeroes of the Bessel function $J_2(\mu T)$, rather than $J_1(\mu T)$. The smoking gun signature of RS1 at the LHC is [9] a tower of resonances with masses 1, 1.83, 2.66, 3.48, ..., in units of the first heavy resonance. For our models, the prediction is a tower of resonances with masses 1, 1.64, 2.26, 2.88, ..., in units of the first heavy resonance.

LHC experiments will also be sensitive to the extra toroidal KK states, if R is of order a TeV. The spectrum of these additional excitations is evenly spaced, distinguishing it from extra toroidal dimensions not associated with AdS directions [8].

LHC experiments may be sensitive to our extra winding states, depending upon which bulk modes have wind-

ing excitations, and how they couple to Standard Model particles. For R of order $1/T$, the winding states have discrete mass spacings of order T . For R of order $1/k$, the winding states are essentially a continuum on top of a mass gap of $5.13T$. This is somewhat reminiscent of LR models [10], but clearly distinct since in that case there is no mass gap in the continuum KK spectrum.

Light winding states have been discussed in the literature [11] in the context of ADD models [12] with a TeV string scale [13]. Here we have obtained TeV scale string winding states without lowering the string scale. This is due to the double warping which is a generic feature of winding states along compactified AdS directions.

Acknowledgments

The authors are grateful to Bill Bardeen, Bogdan Dobrescu, Minjoon Park, and Daniel Robbins for useful discussions. JL acknowledges the support of the Aspen Center for Physics. This research was supported by the U.S. Department of Energy Grants DE-AC02-76CHO3000 and DE-FG02-90ER40560.

-
- [1] L. Randall and R. Sundrum, Phys. Rev. Lett. **83**, 3370 (1999) [arXiv:hep-ph/9905221]; Phys. Rev. Lett. **83**, 4690 (1999) [arXiv:hep-th/9906064].
 - [2] O. Aharony, S. S. Gubser, J. M. Maldacena, H. Ooguri and Y. Oz, Phys. Rept. **323**, 183 (2000) [arXiv:hep-th/9905111].
 - [3] T. Appelquist, B. A. Dobrescu, E. Ponton and H. U. Yee, Phys. Rev. D **65**, 105019 (2002) [arXiv:hep-ph/0201131]; T. Gherghetta and A. Kehagias, Phys. Rev. Lett. **90**, 101601 (2003) [arXiv:hep-th/0211019].
 - [4] T. Gherghetta, E. Roessl and M. E. Shaposhnikov, Phys. Lett. B **491**, 353 (2000) [arXiv:hep-th/0006251].
 - [5] E. Witten, JHEP **9807**, 006 (1998) [arXiv:hep-th/9805112]; M. Alishahiha, arXiv:hep-th/9811042; J. Gomis, A. V. Ramallo, J. Simon and P. K. Townsend, JHEP **9911**, 019 (1999) [arXiv:hep-th/9907022].
 - [6] W. D. Goldberger and M. B. Wise, Phys. Rev. D **60**, 107505 (1999) [arXiv:hep-ph/9907218].
 - [7] É. Mathieu, J. Math. Pures Appl. **13** 137 (1868).
 - [8] H. Davoudiasl, J. L. Hewett and T. G. Rizzo, JHEP **0304**, 001 (2003) [arXiv:hep-ph/0211377].
 - [9] H. Davoudiasl, J. L. Hewett and T. G. Rizzo, Phys. Rev. Lett. **84**, 2080 (2000) [arXiv:hep-ph/9909255].
 - [10] J. Lykken and L. Randall, JHEP **0006**, 014 (2000) [arXiv:hep-th/9908076].
 - [11] A. Donini and S. Rigolin, Nucl. Phys. B **550**, 59 (1999) [arXiv:hep-ph/9901443].
 - [12] N. Arkani-Hamed, S. Dimopoulos and G. R. Dvali, Phys. Lett. B **429**, 263 (1998) [arXiv:hep-ph/9803315].
 - [13] J. D. Lykken, Phys. Rev. D **54**, 3693 (1996) [arXiv:hep-th/9603133].